

On Quantization of the Electrical Charge–Mass

Dmitriy M Palatnik¹

6400 N Sheridan Rd 2605, Chicago, IL 60626

Abstract

Suggested a non-linear, non-gauge invariant model of Maxwell equations, based on the Kaluza-Klein theory. The spectrum of elementary charges and masses is obtained.

PAC numbers: 11.10.-z; 11.10.Lm

1 Introduction

Quantization of the electrical charge and mass still remains a challenge for modern physics. Several attempts were made to solve this problem, e.g. [1, 2]. As an alternative to these consider the following approach. Consider fields $(g_{ab}, \tilde{A}_a)$ in four-dimensional spacetime, where g_{ab} is the metrics and $\tilde{A}_a$ is dimensionless electromagnetic potential, related to the one in standard units by the rule $\tilde{A}_a = \ell A_a$, where ℓ is dimensional constant. Take the following actions for a particle and fields (electromagnetic and gravitational):

$$S_p = -m_1 c_* \int \sqrt{G_{ab} dx^a dx^b} - m_2 c_* \int \tilde{A}_a dx^a; \quad (1)$$

$$S_f = -\frac{1}{16\pi c_* \ell^2} \int d\Omega \sqrt{-g} \tilde{F}^{ab} \tilde{F}_{ab}; \quad (2)$$

$$S_g = -\frac{c_*^3}{16\pi k} \int d\Omega \sqrt{-g} R(g). \quad (3)$$

¹e-mail: palatnik@ripco.com

Here $\tilde{F}_{ab} = \partial_a \tilde{A}_b - \partial_b \tilde{A}_a$, $\tilde{F}^{ab} = G^{ac} G^{bd} \tilde{F}_{cd}$, tensor G^{ab} is inverse to $G_{ab} = g_{ab} - \tilde{A}_a \tilde{A}_b$. The mass of the particle is denoted by m_1 ; ‘mass’ $m_2 = \ell^{-1} c_*^{-2} e$ is related to the electrical charge, e . The speed of light is denoted by c_* , and k is the Newtonian constant of gravitational interactions. This theory may be regarded as a model of the Kaluza-Klein theory. If $\tilde{A}_a$ were a gauge potential and transformations $\tilde{A}_a = \tilde{A}'_a + \partial_a f$ were allowed, one might assume due to (1) that G_{ab} is gauge-invariant. Then, from definition of G_{ab} one would obtain that metrics g_{ab} transforms by the rule $g_{ab} = g'_{ab} + 2\tilde{A}'_{(a} \partial_{b)} f + \partial_a f \partial_b f$. Thus, as it follows from (1)–(3), this theory is *not* gauge-invariant. Define densities, μ_1 and μ_2 , accordingly,

$$m_i = \int \mu_i \sqrt{-g} dV, (i = 1, 2.) \quad (4)$$

Here dV is element of three-dimensional volume. From the least action principle it follows that ‘generalized’ Maxwell equations are:

$$\tilde{F}^{ab}_{;b} + G^{ad} \tilde{F}_{dc} \tilde{F}^{bc} \tilde{A}_b = -4\pi \ell^2 J^a; \quad (5)$$

$$\tilde{F}_{[ab;c]} = 0. \quad (6)$$

Here semicolon denotes the covariant derivative associated with g_{ab} , i.e. equations $g_{ab;c} = 0$ take place. The electrical current,

$$J^a = (\mu_2 c_*^2 - \mu_1 c_*^2 \tilde{A}_b U^b) \frac{U^a}{U^0}. \quad (7)$$

Here

$$U^a = \frac{dx^a}{\sqrt{G_{cd} dx^c dx^d}}. \quad (8)$$

The stress-energy, entering Einstein equations for g_{ab} consists of two parts; T_{ab} , pertaining to the particle, and t_{ab} , pertaining to electromagnetic field,

$$T^{ab} = \mu_1 c_*^2 \frac{U^a U^b}{U^0}; \quad (9)$$

$$t^{ab} = \frac{1}{4\pi\ell^2} \left(-G^{ad} \tilde{F}_{dc} \tilde{F}^{bc} + \frac{1}{4} g^{ab} \tilde{F}_{cd} \tilde{F}^{cd} \right). \quad (10)$$

Below the attention is focused on electromagnetic interactions in flat space-time ($g_{ab} = \eta_{ab} = \text{diag}(1, -1, -1, -1)$).

2 Non-Relativistic Limit

In non-relativistic limit one neglects terms of the order $\frac{v}{C_*}$, where v denotes speeds of particles. Take $x^a = (c_* t, x^\alpha)$, $\alpha = 1, 2, 3$. The potential is described by $\tilde{A}_a(x^b) = (\sin \Psi(x^b), 0, 0, 0)$, generated by the only non-vanishing component of the current,

$$J^0 = \mu_2 c_*^2 - \mu_1 c_*^2 \frac{\tilde{A}_0}{\sqrt{G_{00}}}. \quad (11)$$

Then (5) is equivalent to

$$\Delta \Psi = -4\pi\ell^2 c_*^2 (\mu_2 \cos \Psi - \mu_1 \sin \Psi); \quad (12)$$

here Δ denotes the Laplacian.

One may write down a class of exterior solutions, corresponding to the charge distribution J^0 ,

$$\tilde{A}_0(x^a) = \sin[\ell\phi(x^a) + \psi]. \quad (13)$$

Here $\phi(x^a)$ is the Coulomb potential generated by current component $j^0(x^a) = \cos[\ell\phi(x^a) + \psi]J^0(x^a)$, and ψ is the constant of integration. The exterior solution for a pointlike charge, e , in spherical coordinates, $x^a = (c_*t, r, \theta, \varphi)$,

$$\tilde{A}_0 = \sin\left(\frac{\ell e}{r} + \psi\right). \quad (14)$$

Since field equations are non-linear, the superposition principle doesn't work, and asymptotic on spatial infinity should be the same for any matter distribution.

To find respective 'gravitational' potential G_{ab} and electromagnetic potential, one has to rescale the speed of light. Really, on spatial infinity $ds_1^2 \equiv G_{ab}dx^a dx^b = \cos^2\psi c_*^2 dt^2 - dr^2 - r^2(d\theta^2 + \sin^2\theta d\varphi^2)$, hence, one may introduce observable speed of light, $|c|$, where $c = c_* \cos\psi$; then metrics H_{ab} is defined accordingly, $ds_1^2 = H_{ab}dx^a dx^b$, where $x^a = (ct, x^\alpha)$. In the case considered above,

$$ds_1^2 = \frac{\cos^2\left(\frac{\ell e}{r} + \psi\right)}{\cos^2\psi} c^2 dt^2 - r^2(d\theta^2 + \sin^2\theta d\varphi^2). \quad (15)$$

For 'interval' $ds_2 \equiv -\ell A_a dx^a$ one obtains,

$$ds_2 = -(\cos\psi)^{-1} \tilde{A}_0 c dt - \tilde{A}_\alpha dx^\alpha, \quad (16)$$

where A_a is electromagnetic potential in the standard units. In considered above spherically-symmetric case,

$$A_0(r) = \frac{\sin\left(\frac{\ell e}{r} + \psi\right)}{\ell \cos\psi}. \quad (17)$$

For $\ell e \ll r$ one obtains from (15) and (17),

$$H_{00} = 1 - \frac{2\ell e \tan \psi}{r} + O(r^{-2}); \quad (18)$$

$$A_0 = \ell^{-1} \tan \psi + \frac{e}{r} + O(r^{-2}). \quad (19)$$

3 Quantizing the Electrical Charge

Consider flat spacetime with motionless electrical charge, described in spherical coordinate system. Take $\tilde{A}_a = [\sin \frac{\Phi(r)}{r}, 0, 0, 0]$. Equation (5) is equivalent to the following,

$$\Phi'' = 4\pi\ell^2 c_*^2 r \left(-\mu_2 \cos \frac{\Phi}{r} + \mu_1 \sin \frac{\Phi}{r} \right); \quad (20)$$

here prime denotes differentiation over r .

One may take the following model of the current (7) for the particle. (a)

For the density take

$$-\mu_1 + i\mu_2 = \frac{p^2}{4\pi\ell^2 c_*^2} \exp(i\psi) \frac{\arctan \sqrt{\tilde{A}_a \tilde{A}^a} + \psi}{\sin \left(\arctan \sqrt{\tilde{A}_a \tilde{A}^a} + \psi \right)}. \quad (21)$$

Here $\tilde{A}^a = G^{ab} \tilde{A}_b$, and p, ψ are real normalizing constants, found from (4).

(b) For four velocity U^a take

$$U^a = \frac{\tilde{A}^a}{\sqrt{\tilde{A}_b \tilde{A}^b}}. \quad (22)$$

(c) Define boundary, S , of the particle by equation,

$$\left(\arctan \sqrt{\tilde{A}_a \tilde{A}^a} + \psi \right)_S = 0. \quad (23)$$

It is assumed that in the exterior of the boundary $\mu_1 + i\mu_2 = 0$.

Solving (20) with (21), one obtains, $\tilde{A}_- = \sin\left(C\frac{\sin(pr)}{pr} - \psi\right)$ and $\tilde{A}_+ = \sin\left(\frac{a}{r} + b\right)$, respectively interior and exterior solutions. Here C , a , and b are constants of integration.

Define boundary of the particle, as a surface of the sphere $r = R$. Then, from (23) it follows, (a) $\sin(pR) = 0$. Demanding continuity of the potential together with its first derivative, one obtains two relations: (b) $b = -\frac{a}{R} - \psi$; (c) $a = -CR\cos(pR)$. From (a) it follows that two cases take place: (i) $a = -C\frac{2\pi n}{p}$ with $R = \frac{2\pi n}{p}$ and (ii) $a = C\frac{\pi(2n+1)}{p}$ with $R = \frac{\pi(2n+1)}{p}$. In both cases $n = 0, 1, 2, \dots$. For the exterior potential one obtains two branches, corresponding to cases (i) and (ii) above. Assuming that both branches have the same asymptotic on spatial infinity, one obtains, $C = \pi N$, $N = 0, \pm 1, \pm 2, \dots$. On the other hand, demanding that mass-charge be finite, one has to put $N = \pm 1$. Thus, in the standard units, introducing e , charge of proton, one obtains,

$$A_{0(1)} = \ell^{-1} \cos^{-1} \psi \sin\left(N\frac{(2n)\ell e}{3r} + \psi\right); \quad (24)$$

$$A_{0(2)} = \ell^{-1} \cos^{-1} \psi \sin\left(-N\frac{(2n+1)\ell e}{3r} + \psi\right). \quad (25)$$

Both branches originate from the following interior ($r \leq R$) solution:

$$A_{0(-)} = \ell^{-1} \cos^{-1} \psi \sin\left(\pi N\frac{\sin(pr)}{pr} - \psi\right). \quad (26)$$

Here $p = 3\pi^2\ell^{-1}e^{-1}$. The respective mass-charge density is

$$\mu_1 - i\mu_2 = -\frac{p^2 \exp(i\psi)}{4\ell^2 c_*^2} \frac{v(pr)}{\sin(\pi v(pr))}, \quad (27)$$

and it doesn't depend on N . Thus, one may take $N = 1$. In (27) $v(x) = \frac{\sin x}{x}$.

Expanding solutions (24) and (25), one obtains,

$$A_{0(1)} = A_* + \frac{e}{3} \left(\frac{2n}{r} \right) - \frac{s}{2} \left(\frac{2n}{r} \right)^2 - \frac{w}{6} \left(\frac{2n}{r} \right)^3 + \dots \quad (28)$$

$$A_{0(2)} = A_* - \frac{e}{3} \left(\frac{2n+1}{r} \right) - \frac{s}{2} \left(\frac{2n+1}{r} \right)^2 + \frac{w}{6} \left(\frac{2n+1}{r} \right)^3 + \dots \quad (29)$$

Here $A_* = \ell^{-1} \tan \psi$, and the following definitions are used for charges,

$$e = 3\pi^2 \ell^{-1} p^{-1}; \quad (30)$$

$$s = \pi^4 \ell^{-1} p^{-2} \tan \psi; \quad (31)$$

$$w = \pi^6 \ell^{-1} p^{-3}; \quad (32)$$

or, expressing p, ψ , and ℓ through e, s , and w , one obtains,

$$p^2 = \frac{\pi^4}{3} \frac{e}{w}; \quad (33)$$

$$\tan \psi = \frac{\sqrt{3}s}{\sqrt{ew}}; \quad (34)$$

$$\ell = \frac{3\sqrt{3}}{e} \sqrt{\frac{w}{e}}. \quad (35)$$

The spectrum of charges consists of two parts; the first one is $q_{2n} = \frac{2n}{3}e$, ($n = 1, 2, \dots$), and the second one is $q_{2n'+1} = -\frac{2n'+1}{3}e$ ($n' = 0, 1, 2, \dots$), these correspond to quarks and leptons.

4 Quantizing the Mass

Insofar, parameter ψ isn't fixed yet. To fix it, one may use (4) together with quantization rule for charge. From (4) and (27) it follows

$$m_{1(k)} - im_{2(k)} = -\frac{\pi \exp(i\psi)}{p\ell^2 c_*^2} \alpha_k; \quad (36)$$

$$\alpha_k = \int_0^{\pi k} \frac{x^2 v(x) dx}{\sin(\pi v(x))}. \quad (37)$$

Here $k = 2n$ for electrical charge $\frac{2n}{3}e$, and $k = 2n + 1$ for electrical charge $-\frac{2n+1}{3}e$. On the other hand, $m_{2(2n)} = -\frac{\pi^2(2n)}{p\ell^2 c_*^2}$, and $m_{2(2n+1)} = \frac{\pi^2(2n+1)}{p\ell^2 c_*^2}$. It follows, then, $\sin \psi_{2n} = -\frac{\pi(2n)}{\alpha_{2n}}$, and $\sin \psi_{2n+1} = \frac{\pi(2n+1)}{\alpha_{2n+1}}$. Hence, one obtains spectrum of masses,

$$m_{2n} = \frac{e}{3\pi\ell c_*^2} \sqrt{\alpha_{2n}^2 - 4\pi^2 n^2}; \quad n = 1, 2, \dots \quad (38)$$

$$m_{2n'+1} = \frac{e}{3\pi\ell c_*^2} \sqrt{\alpha_{2n'+1}^2 - \pi^2(2n'+1)^2}; \quad n' = 0, 1, 2, \dots \quad (39)$$

here masses m_{2n} correspond to charges q_{2n} , and masses $m_{2n'+1}$ correspond to charges $q_{2n'+1}$. For the first three alphas one obtains, $\alpha_1 \approx 6.27$; $\alpha_2 \approx 30.2$; $\alpha_3 \approx 93.5$. The respective masses for two quarks are $m_{-\frac{1}{3}} = .059m_e$ and $m_{\frac{2}{3}} = .32m_e$, where m_e is mass of an electron.

5 Conclusion

In this work we obtained spectrum of electrical charges, corresponding to leptons, (e, μ, τ) , and quarks (u, d, s, c, t, b) . It appears, that spectrum of masses (38) and (39) doesn't describe quarks (s, c, t, b) . On the other hand, correct spectrum of electrical charges allows to believe that the analogous result may follow from the Kaluza-Klein theory without assumption of topological closeness of the extra-dimension.

Definition of density (21) is made accordingly in order to obtain the simplest differential equation for potential (20). A more consistent and realistic

approach would be considering constant densities, μ_1 and μ_2 . The price to be paid for that is to deal with non-linear equation for potential. We thought of doing that as unworthy, since the theory anyway is just a model of Kaluza-Klein theory, so that the effort of solving non-linear equation for potential should be made in frames of Kaluza-Klein.

The masses are calculated for isolated particles. In realistic situations of many interacting particles constant ψ should be found for the complete closed system which would shift the masses (but not the electrical charges).

6 Acknowledgments

I wish to thank Boris S Tsirelson for helping me with numeric computations.

References

- [1] P.A.M. Dirac, *Phys. Rev.* **74**, 817 (1948).
- [2] H. Georgi, S.L. Glashow *Phys. Rev. Lett.* **32**, 438 (1974).